FOR EMERGENCIES/EXTREME STRESS SITUATIONS (TO AVOID OR CORRECT ADDISONIAN CRISIS)

Conditions which must be treated immediately:

- severe injury of any type
- blood loss
- fluid and/or electrolyte loss
- infection
- severe vomiting

1.) I.V. hydrocortisone.

2.) Administer I.V. isotonic saline.

· diarrhea.

Treatment protocol prior to Emergency Room arrival:

- Give Hydrocortisone 100 mg. IM (intramuscularly) in thigh or gluteus muscle.
- Patient then should be immediately admitted to the nearest Emergency Room, where adrenal crisis treatment should begin promptly.
- If transportation to emergency medical facility is impossible, treat patient on site as listed below.

Protocol for treatment of adrenal crisis in emergency room:

3.) Continue I.V. hydrocortisone until oral medication is tolerated.

| For further details, call Dr Specialty: Office Phone: Emergency Pager or Exchange #: Address | | |
|--|-----------------|----------|
| City | State | Zip |
| Patient NameAddress | Home Phone: | |
| | | Zip |
| MEDICAL CONDITIONS | MEDICATION | DOSAGES |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Emergency | Relationship to | patient: |
| Home phone: | | |
| Emergency contact #2: | Relationship to | patient: |
| | | Cell: |